



Allergies (also specify reaction) None known

Date

(YYYY MM DD)

Write firmly for legible copy

**Emergency Department
Symptoms Suggestive of Hip Fracture**

Form D648 (Rev. 09.2011)

Item	Order	Transcribed
1	<p>Clinical Assessment Patient presents with symptoms suggestive of hip fracture:</p> <ul style="list-style-type: none"> • Pain in the hip, groin and/or knee • Severe pain with movement of affected limb **Assess verbal and non-verbal pain cues** • External rotation and/or shortening of affected limb • Inability to weight bear on affected side 	
2	<p>Clinical Diagnostics</p> <p><input type="checkbox"/> CBC, Electrolytes, Urea, Creatinine, Blood glucose, PT, INR, Group and Screen</p> <p><input type="checkbox"/> ECG</p> <p><input type="checkbox"/> X-ray of affected hip: <input type="checkbox"/> Right hip <input type="checkbox"/> Left Hip</p> <p><input type="checkbox"/> X-ray – Other _____</p> <p><input type="checkbox"/> Additional Diagnostics: _____</p>	
3	<p>Clinical Interventions</p> <p><input type="checkbox"/> Keep patient NPO</p> <p><input type="checkbox"/> Initiate IV therapy with 0.9% normal saline. Bolus _____ mL over _____ minutes.</p> <p><input type="checkbox"/> Initiate IV therapy with 0.9% normal saline @ _____ mL/hr</p> <p><input type="checkbox"/> Insert urinary catheter to straight drain for female patient</p>	
4	<p>Clinical Monitoring</p> <ul style="list-style-type: none"> • Document temperature, heart rate, respiratory rate, oxygen saturation, blood pressure and pain scale prior to and 30 min to 1 hour after administration of analgesia • ED MD to consider use of Iliofascial 3 in 1 block to reduce the need for systemic analgesic. Monitor effect of Iliofascial block • Document neurovascular status of affected limb (using 5P's: pain, pulses, pallor, paresthesia, paralysis) Q4H • Document ISAR score if patient >65 years 	
5	<p>Medications</p> <p><i>*For patients with known Dementia, consider beginning with HYDROmorphine (Dilaudid) 0.5mg IV</i></p> <p><input type="checkbox"/> HYDROmorphine (Dilaudid) 0.5 -1mg IV Q1hour PRN until pain relieved. If pt requires 4mg in 4 hours, please contact MD to reassess.</p> <p>_____</p> <p><input type="checkbox"/> Ondansetron 4mg IV Q8H PRN for nausea and vomiting</p> <p>_____</p> <p>Other _____</p>	
6	<p>Consults</p> <p><input type="checkbox"/> Orthopedics <input type="checkbox"/> Anesthesia <input type="checkbox"/> General Medicine <input type="checkbox"/> Geriatrics <input type="checkbox"/> Social Work <input type="checkbox"/> Pharmacy</p>	

Date	Time	Print Name	Signatures
(YYYY MM DD)	(HH : MM)	_____	_____, M.D.
(YYYY MM DD)	(HH : MM)	_____	_____, R.N.

